

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.

1.9
F7626R
Exp. 2

U. S. DEPT. OF AGRICULTURE
LIBRARY
AUG 24 1961
CURRENT SEWAGE

FOREST SERVICE - U. S. DEPARTMENT OF AGRICULTURE

PACIFIC SOUTHWEST
FOREST AND RANGE
EXPERIMENT STATION
BERKELEY - CALIFORNIA

RESEARCH NOTE

No. 178

April 1961

GIBBERELLIN AND THIOUREA BREAK SEED DORMANCY IN CALIFORNIA CEANOTHUS

Lowell Adams, Wildlife Research Biologist
Eugene Stefanescu, Range Aid (Research)
David J. Dunaway, Range Aid (Research)

Division of Range Management and
Wildlife Habitat Research

Seeds of most wild shrubs do not germinate until they are "winterized" naturally or by artificial treatment. Consequently, seeding of shrubs for game range improvement is nearly always faced with this immediate handicap. One of the most common methods of breaking seed dormancy is stratification in cold moist sand. This takes up to three months, however, and the seeds must be kept moist thereafter until planted. These requirements present cumbersome limitations in large-scale, practical seedings. In recent years considerable progress has been made in breaking dormancy with chemicals.

This report describes a newly-discovered chemical treatment for breaking seed dormancy in three species of *Ceanothus*: mountain whitethorn (*C. cordulatus* Kell.), Lemmons ceanothus (*C. lemmonii* Parry), and buckbrush ceanothus (*C. cuneatus* (Hook)). The chemicals used are gibberellin and thiourea.

The treatment that increased germination most consisted of:

- (1) Soaking seeds in boiling water one minute,
- (2) Soaking in gibberellin (400 parts per million) 13 hours,
- (3) Drying the seeds 4 days, and
- (4) Soaking in 3 percent thiourea five minutes to one hour.

Variations in soaking times and chemical concentrations gave graded responses less effective than the above procedure.

A variety of treatments was applied to four other species without success:

<u>Species</u>	<u>Treatments</u>
Bigberry manzanita (<u>Arctostaphylos glauca</u> Lindl.) and bittercherry (<u>Prunus emarginata</u> (Dougl.) Walp.)	Same gibberellin-thiourea treatments given successfully to the three ceanothus species.
Deerbrush (<u>Ceanothus integerrimus</u> H. & A.) and Fremont silktassel (<u>Garrya Fremontii</u> Torr.)	A great variety including various combinations and levels of hot water, thiourea, gibberellin, heat, cold and light.

These species and treatments are not included in the following description of the successful experiments, but information will be supplied upon inquiry to the authors.

The Experiments and Results

We submitted the seeds of the three species of ceanothus first to treatments for dormancy attributable to the seed coat, and second to treatments for dormancy in the seed itself.

Seed-coat Treatments

A seed lot from each species was immersed in each of the following chemicals that have been used by others to make seed coats permeable:

<u>Chemical</u>	<u>Concentration</u> (Percent)	<u>Soaking Time</u> (Hours)
sodium hypochlorite	5.25	1/2, 2, 5, 24
acetone	100.00	1/2, 2, 5, 24
xylol	100.00	1/2, 2, 5, 24
methyl alcohol	95.00	1/2, 2, 5, 24
ether	100.00	1/2, 2, 5, 24
sulfuric acid	conc.	1/2, 1 1/2, 2 1/2
hot water		steep or boil 1/2

1/2 Steep: seeds placed in one liter of water at 180°F. and allowed to cool to room temperature. Boil: seeds immersed in boiling water 1, 4 or 8 minutes.

The test of effectiveness of seed-coat treatments was whether the treated seeds plumped (imbibed water) after treatment when placed on wet blotting paper in petri dishes. One set of untreated seeds was placed in germination dishes as controls.

The seeds of all three ceanothus species plumped after all treatments at all levels. With no treatment, a small percentage of the buckbrush seeds, but no others, plumped.

Seed Treatments

Since all seed-coat treatments resulted in plumping in ceanothus, we used the simplest treatment--boiling for one minute--to induce seed-coat permeability before seed treatment. Then seeds were immersed 13 hours in gibberellin (gibberellic acid neutralized with potassium hydroxide) at three concentrations: 200, 400, and 800 parts per million. A fourth lot of seeds received no gibberellin treatment.

After drying for 4 days, each lot of gibberellin-treated seed, plus an untreated lot, was immersed in 3 percent thiourea for 5 minutes and for 1, 24, and 72 hours. Thus, individual lots of the seeds of each species of shrub were subjected to one of 16 treatments (fig. 1).

Thiourea without gibberellin had practically no effect except in buckbrush, where we achieved about the same results as Hubbard 1/ who used similar treatment. The gibberellin more than doubled the germination rate over the treatment with thiourea alone.

We found no deformed seedlings such as McConnell 2/ reported in gibberellic acid-treated bitterbrush.

Since varietal or strain differences may affect response to dormancy treatments, the sources of seeds we used are listed below. All are from California.

Manzanita - Paradise, Butte County
Cherry - Snag Lake, Butte County
Deerbrush - "Southern California"
Mountain whitethorn - Yellow Creek, Plumas County
Lemmons ceanothus - Magalia, Butte County
Buckbrush, Paradise, Butte County
Silktassel - "Near Paradise"

All seeds were purchased from commercial seed sources and were tested within 1 year after purchase.

1/ Hubbard, Richard L. Hot water bath and thiourea break dormancy of wedgeleaf ceanothus seed. U. S. Forest Serv. Calif. Forest and Range Expt. Sta. Res. Note 143. 4 pp. 1958.

2/ McConnell, Burt R. Effect of gibberellic acid and cold treatments on the germination of bitterbrush seed. U. S. Forest Serv. Pacific Northwest Forest and Range Expt. Sta. Res. Note 187. 4 pp. 1960.




Figure 1.--Numbers of seeds germinated in 50-seed lots treated with four levels of gibberellin and four of thiourea.

